Malaysian Organ Sharing System (MOSS)

Editor - The extensive media coverage of organ transplantation over the last two years in this country has resulted in an increase in the number of cadaveric transplantation. There is now a growing concern about how cadaveric organs are allocated and to whom; with particular reference to distributive justice and medical suitability.

The Malaysian Society of Nephrology and Ministry of Health of Malaysia have initiated the development of a national organ sharing network and is named Malaysian Organ Sharing System (MOSS). Presently MOSS is only involved in the allocation of cadaveric kidneys as the number of patients in the waiting list for other solid organs are still very small. MOSS had developed a system of organ (cadeveric kidneys) allocation based on distributive justice and medical suitability and is described below.

Eligibility

All Malaysian citizens who are medically fit for transplantation are eligible to be registered with MOSS as potential cadaveric renal recipient^{1,2}. Permanent residents will be considered on an individual basis.

Exclusion Criteria

Patients aged less than 2 years old or above 60 years old are disqualified^{3,4}. Patients with any disease or illness with expected survival of less than 5 years or with a resultant poor quality of life are not eligible. This will include patients with:

- 1. Positive HIV serology⁵.
- 2. Positive Hepatitis Be antigen⁶.
- 3. Active liver disease^{6,7}.
- 4. Severe cardiovascular/ cerebrovascular/ pulmonary disease.
- Dementia or psychosis with no underlying treatable disease.

- 6. Malignancy of limited life expectancy^{8,9}.
- 7. Oxalosis (not contraindicated for combined liver and kidney transplantation).
- 8. Fabry's disease.
- 9. Active substance abuse.

Pre-registration Assessment

Cardiac assessment (including cardiac stress test & echocardiogram)^{3,10} is required for asymptomatic, nondiabetic patients ≥55 years old or diabetic patients ≥35 years old. For symptomatic patients or asymptomatic patients with positive stress test, they should be further assessed by a cardiologist and treated appropriately.

Criteria for Prioritization of Candidates 11-18

With rare exception, the donor and recipient should be of the same ABO blood group type. One of the two kidneys will also be allocated to the procurement centre.

Candidates with limited life expectancy (<1 year) without renal transplant will be placed into medical emergency list (SOS list). This decision can only be made by a nephrologist and a committee will review the SOS list periodically. Candidates on the SOS list will receive top priority in receiving a cadaveric kidney.

Candidates will be selected based on the following scoring system:

- A. HLA Ag (A,B and DR) matching *** (12 points) ***Six HLA antigens match will be given top priority.
- ***Due to logistic reasons, HLA Ag matching is presently not being used as a selection criteria.

LETTERS TO EDITOR

- B. PRA (10 points)
- C. Waiting time (20 points)
- D. Logistical Score (6 points)
- E. Age less than 18 years

Patients less than 18 years old will be given extra weightage.

H S Wong, MRCP Hononary Secretary Malaysian Organ Sharing System (MOSS) Malaysian Society of Nephorology (MSN)

References

- Kleinig J. Organ Transplants: The just and human expectations of nonresidents. Transplantation Proceedings 1989; 21: 3430-1.
- Prottas JM. Nonresident aliens and access to organ transplant. Transplantation proceedings 1989; 21: 3426-9.
- Ismail N, Hakim RM, Helderman JH. Renal replacement therapies in the elderly: Part II. Renal transplantation AJKD 1994; 23: 1-15.
- De Shazo CV, Simmons RL, Bernstein DM et al. Results of renal transplant in 100 children. Surgery 1974; 76: 461.
- Allen RDM, Chapman JR. The Recipient. In: A manual of renal transplantation. Little, Brown and Company, 1994: 8-20.
- Durlik M, Gaciong Z, Soluch L et al. Effect of chronic liver disease on clinical outcome in renal allograft recipients. Transplantation Proceedings 1997; 29: 787-90.
- Goffin E, Pirson Y, Cornu C et al. Outcome of HCV infection after renal transplantation. Kid Int 1994; 45: 551-5.
- 8. Penn I. Cancer in the immunosuppressed organ recipient. Transplantation Proceeding 1991; 23: 1771-2.
- Penn I. The effect of immunosuppression on pre-existing cancers. Transplantation 1993; 55: 742-7.
- Logue RB. Evaluation and management of patients with heart disease who undergo non cardiac surgery. In: JW Hurst (6th ed). The Heart. McGraw-Hill Book Company 1986; 1511-9.

- 11. Najarian JS, Howard RJ, Foker JE, Simmons RL. Renal transplantation: criteria for selection and evaluation of patients and immunological aspects of transplantation. In Brenner BM, Rector FC. The Kidney. WB Saunders Company 1976: 1745-95.
- Starzl TE, Hakala TR, Tzakis A et al. A multifactorial system for equitable selection of cadaver kidney recipients. JAMA 1987; 257: 3073-5.
- 13. Starzl TE, Shapiro R, Tepermen L The point system for organ distribution. Transplantation Proceeding 1989; 21: 3432-6.
- 14. Lazda VA, Blaesing ME. Is allocation of kidneys on basis of HLA match equitable in multiracial populations? Transplantation Proceedings 1989; 21: 1415-6.
- 15. Hata Y, Cecka JM, Takemoto S *et al*. Effects of changes in the criteria of nationally shared kidney transplants for HLA-matched patients. Transplantation 1998; 208-12.
- Gjerston DW, Terasaki PI, Takemoto S et al. National allocation of cadaveric kidneys by HLA matchingprojectd effect on outcome and costs. N Engl J med 1991; 324: 1032-6.
- Wujciak T, Opelz G. Computer analysis of cadaver kidney allocation procedures. Transplantation 1993; 55: 516-21.
- Greenstein SM, Schechner RS, Louis P et al. Evidence that zero antigen-matched cyclosporine treated renal transplant recipients have graft survival equal to that of matched recipients - reevaluation of points. Transplantation 1990; 49: 332-6.